

Indiana University - Purdue University Fort Wayne Opus: Research & Creativity at IPFW

2012 IPFW Student Research and Creative
Endeavor Symposium

IPFW Student Research and Creative Endeavor
Symposium

4-14-2012

Release and Recovery of Trapped Fullerenes through the Use of Organic Acids

Amanda Sauders

Indiana University - Purdue University Fort Wayne

Follow this and additional works at: http://opus.ipfw.edu/stu_symp2012



Part of the [Chemistry Commons](#)

Recommended Citation

Amanda Sauders (2012). *Release and Recovery of Trapped Fullerenes through the Use of Organic Acids*.
http://opus.ipfw.edu/stu_symp2012/10

This Presentation is brought to you for free and open access by the IPFW Student Research and Creative Endeavor Symposium at Opus: Research & Creativity at IPFW. It has been accepted for inclusion in 2012 IPFW Student Research and Creative Endeavor Symposium by an authorized administrator of Opus: Research & Creativity at IPFW. For more information, please contact admin@lib.ipfw.edu.

“Release and Recovery of Trapped Fullerenes through the Use of Organic Acids”

Amanda Sauders
Dr. Steven Stevenson
Chemistry Department
Indiana University-Purdue University Fort Wayne

Samples of the recently discovered metallic nitride fullerenes (MNFs) are in high demand for application development due to their interesting chemical and physical properties. The "Stir and Filter Approach" (SAFA) effectively purifies the most chemically inert MNF (e.g., Sc₃N@Ih-C₈₀) due to its lack of reactivity with diaminosilica, the separation medium. The problem, to date, with this separation method is an irreversible binding of empty-cage fullerenes (e.g., C₆₀, C₇₀, C₈₄) and the more reactive MNFs (e.g., Sc₃N@C₆₈, Sc₃N@C₇₈, Sc₃N@D_{5h}-C₈₀) to the silica. In lieu of discarding the spent SAFA silica to waste, we have instead developed a novel method for releasing the trapped fullerenes. The recovered fullerenes can then be re-used in subsequent purification methods for the isolation of more MNF samples. In this presentation, we will discuss specific procedures for removing the fullerenes from the spent silica which includes the utilization of an array of organic acids. We will also describe the selective release chemistry based on the effect of the different acids.